

ASL-contrast-free acquisition technique is ideal for patients in whom contrast is contraindicated. It gives a simple and fast automated image post processing. 3D ASL helps to rule out focal or global perfusion defects, in geriatric as well as pediatric patients. ASL generates excellent image quality with whole brain coverage

A case of Clinical Benefits of Arterial Spin Labeling (ASL)

Dr. O. P. Gupta

Director

Dr. O. P. Gupta Imaging Centre

Meerut, India



Dr. Muhammad Qasim

Chief Reporting Radiologist

Dr. O. P. Gupta Imaging Centre

Meerut, India



"Arterial Spin Labeling (ASL) is an effective and non invasive technique for follow up of tumors in the brain"

Patient history

- A 55-year-old female
- Recently operated case of Haemangio-Pericytoma grade II tumor
- Right Parieto-occipital craniotomy and complete tumor resection done

Technique & findings


A small post surgical cavity in right occipital region with foci of blood products at periphery and mild surrounding edema with minimal peripheral enhancement were seen. ASL showed low CBF in the enhancing small nodule at the periphery of the surgical cavity suggesting hypoperfusion of inflammatory enhancement and not the residual tumor.

Acquisitions Parameters: ASL


Imaging Mode:	3D
FOV:	24cm
Slice Thickness:	4 mm
Frequency:	512
Phase:	8
Band width:	62.50 Nex-2
Location perslab:	30




Routine MR view




AX DWI



AX T2



AX GRE



AX T1

ASL findings:

Parenchyma Anterior to Post Surgical Cavity

Parameter	ROI	ROI	% of parameter
	(In the parenchyma surrounding surgical cavity)	(contra lateral normal region)	(around cavity as compared to normal region)
CBF	57.850	62.440	92%



Parenchyma Medial to Post Surgical Cavity

Parameter	ROI	ROI	% of parameter
	(In the parenchyma surrounding surgical cavity)	(contra lateral normal region)	(around cavity as compared to normal region)
CBF	34.640	66.560	52%



Enhancing Nodule Posterior to Post Surgical Cavity

Parameter	ROI	ROI	% of parameter
	(In the parenchyma surrounding surgical cavity)	(contra lateral normal region)	(around cavity as compared to normal region)
CBF	26.567	80.267	33%



Parenchyma Anterior to Post Surgical Cavity



Parenchyma Medial to Post Surgical Cavity



Enhancing Nodule Posterior to Post Surgical Cavity



Conclusion

ASL is an excellent technique for diagnosis and follow-up of tumors within the brain.

Image courtesy to Dr. O. P. Gupta Imaging Centre

"This article has been published by GE Healthcare with consent of the authors/institutions concerned in academic interest. The views expressed in the case study are solely those of the Author. For circulation within India only"

The editor sincerely acknowledges the support and contribution of Mr. Vignesh Singh, Senior Application Specialist, India at GE Healthcare.

To learn more about GE MR Systems SMS **GEHC** to **56677** or call our toll free number at **1800 209 9003** or email to ajithkumarDR@ge.com/dhrumil.sorathia@ge.com